

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau




(43) International Publication Date  
31 December 2003 (31.12.2003)

PCT

(10) International Publication Number  
**WO 2004/000037 A1**

- (51) International Patent Classification<sup>7</sup>: **A23K 1/16**
- (21) International Application Number:  
PCT/FI2003/000504
- (22) International Filing Date: 19 June 2003 (19.06.2003)
- (25) Filing Language: Finnish
- (26) Publication Language: English
- (30) Priority Data:  
20021234 24 June 2002 (24.06.2002) FI
- (71) Applicant (for all designated States except US): **KEMIRA CHEMICALS OY** [FI/FI]; P.O. Box 330, FIN-00101 Helsinki (FI).
- (72) Inventors; and  
(75) Inventors/Applicants (for US only): **MIETTINEN, Harri** [FI/FI]; Wiikintie 4 A 3, FIN-02400 Kirkkonummi (FI). **LÖFGREN, Timo** [FI/FI]; Hannikka 32 A, FIN-02360 Espoo (FI).
- (74) Agent: **BERGGREN OY AB**; P.O. BOX 16, FIN-00101 Helsinki (FI).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GROWTH PROMOTOR



(57) Abstract: The invention relates to an animal growth promoting composition. The liquid solution comprises 1.1 to 5 % by weight of sorbic acid or a sorbate calculated as sorbic acid, 45 to 93.9 % by weight of formic acid and 5 to 50 % by weight of propionic acid. The invention also relates to a method for preparing the composition and to its use.

## Growth promotor

5 The invention relates to a composition for promoting the growth of animals and to a method for preparing and using the composition.

Antibiotics have been added to the feed of slaughter animals in order to prevent diseases and to increase meat production. The use of antibiotics may result in resistant bacterial strains, which risk to be carried to humans along with the meat.  
10 This is the reason for efforts made to replace antibiotics with other compounds.

Antibiotics have lately been replaced with organic acids. Thus, for instance, the increase in weight of slaughter animals has been enhanced by means of sorbic acid (*Kirchgessner M. et. al.*; *Journal of Animal Physiology and Animal Nutrition* 74  
15 (1995), 235-242). However, sorbic acid has proved too expensive to be adopted as an alternative as such.

Studies have also confirmed that formic acid acts as a growth promotor in the feed of weaned pigs and swine grown for pork meat. It has also been found that the use  
20 of a mixture of formic acid and potassium sorbate results in better efficiency compared to the use of formic acid (*Partanen K. et. al.*; *Livestock Production Science* 73 (2002) 139-152).

The combined use of formic acid and potassium sorbate involves the problem of  
25 one of the components being liquid and the other solid. It would be desirable to find a means of providing a liquid product containing formic acid and sorbic acid or a sorbate that can be dispensed in liquid state also at low temperatures. Sorbate and sorbic acid are so poorly soluble in formic acid that an efficient amount cannot be dissolved.

30 The use of sorbic acid and its salts as food and feed preservatives is also known. Sorbate in acid form alone is known to possess anti-microbial properties. Consequently, the effect of salts has been enhanced by adding an acid besides sorbate to the material to be preserved, as described in EP patent application 0275  
35 958, among other publications.

Chinese patent application CN 1 269 979 discloses an anti-mildew feed preservative based on potassium sorbate, which has been boosted with an organic acid. The organic acid may be formic acid, propionic acid or a mixture of these. Sorbate crystals are milled to a 60 to 150 mesh grinding fineness, the acid is added and the mixture is allowed to react completely at constant temperature. The mixture ratio has been selected so as to generate a solid reaction product during the reaction. When the solid product thus obtained is crushed, the desired preservative is obtained.

- 10 The solutions of US patent specification 4,083,999, containing 0.6 to 20% of sorbic acid in propionic acid, are previously known in an anti-mildew method for preserving animal feed.

15 The purpose of this invention is to provide a liquid product, which can be dispensed in liquid state also at low temperatures, while still utilising the animal growth promoting effects especially of formic acid and sorbate. As a solution to this problem, liquid compositions were found, which contain both sorbic acid or a sorbate and formic acid as a homogenous solution. These purposes can be surprisingly achieved with the aid of a composition of formic acid and sorbic acid or a sorbate and propionic acid. Calculated on their weight, the compositions of the invention contain

- 1.1 to 5% of sorbic acid or a sorbate calculated as sorbic acid,
- 45 to 93.9% of formic acid and
- 25 - 5 to 50% of propionic acid.

The composition of the invention contains preferably 2 to 5 % by weight of sorbic acid or sorbate calculated as sorbic acid, 45 to 90 % by weight of formic acid and 8 to 50 % by weight of propionic acid.

30

The sorbate is preferably potassium sorbate.

The concentration of formic acid is preferably at least 70 % by weight.

- 35 In preferred compositions, a portion of the acid content of the composition is neutralised. In a preferred composition, the formic acid is partly neutralised with ammonia.

The composition of the invention may further contain water.

5 A preferred composition of the invention contains 2 to 4 % by weight of sorbic acid or a sorbate calculated as sorbic acid, 70 to 90 % by weight of partly neutralised formic acid and 8 to 20 % by weight of propionic acid and 0 to 20 % by weight of water.

10 In accordance with the invention, the composition can be prepared by dissolving sorbic acid or a sorbate in propionic acid, thus yielding soluble sorbic acid or sorbate. Formic acid is then added to this solution, thus forming a solution ready for use.

15 The characterising features of the invention are described in the accompanying claims 1 to 16.

20 It has been found that antibiotic feed additives can be replaced in the feed of slaughter animals with a growth-promoting composition, which contains sorbic acid, formic acid and propionic acid. The anti-microbial and animal growth promoting effect of sorbic acid or sorbate has been enhanced by using acid conditions. Due to the high cost of sorbic acid or sorbate, it was desirable to find an economical, but still efficient active acid for use with sorbic acid or sorbate. Formic acid as such has proved an efficient growth promoter with regard to slaughter animals. It has proved to have an advantageous effect especially in the feeding of  
25 pigs and swine grown for pork meat in reducing the tendency of swine to develop diarrhoea and coliform bacteria in the alimentary canals. As the poor solubility of sorbic acid or sorbate in formic acid was resolved by first dissolving it in propionic acid and the actual solution ready for use was obtained by adding the solution of sorbic acid/sorbate/propionic acid to formic acid, the liquid composition of the  
30 invention was obtained, which proved a highly efficient growth promoter for slaughter animals.

During the use of the product of the invention, it was noted that slaughter animals, such as beef cattle and especially swine developed less diarrhoea and that their meat  
35 mass increased appreciably.

The preparation of the invention can be administered in connection with the feeding of slaughter animals by adding it to the feed. The growth promoter can be added to the feed of slaughter animals either in liquid form or, if the circumstances so require, absorbed into a solid carrier.

5

The feed may be any organic product suitable for slaughter animals, such as feed raw materials, feed preparations, corn, vegetable or organic protein feed. The feed proper is fed to the animals in liquid form admixed in water or in solid form.

10 The following examples are intended to illustrate the invention, yet without limiting the invention to the accompanying examples.

The accompanying figure 1 illustrates the solubility curves of potassium sorbate at different temperatures.

15

### Examples

In the following examples, sorbic acid was added in the form of potassium salt in all of the examples. However, in the presence of formic acid (pK 3.73), potassium sorbate occurs primarily as sorbic acid in the final solution, because the pK value of sorbic acid is 4.76, and correspondingly, the pK value of propionic acid is 4.86.

20

The partly neutralised formic acid product used in the examples contains 76 % by weight of formic acid, 5.5 % by weight of ammonium formate and 18.5 % by weight of water.

25

### Examples 1 to 4 (comparison)

Potassium sorbate in varied amounts was dissolved in a partly neutralised formic acid product by heating so that clear solutions were obtained. Samples of these were stored at room temperature (about 22 °C), 4 °C and – 18 °C. At the end of about 10 days, the crystallisations formed in the samples were examined, with the results shown in the table. In an examination made at the end of about 50 days, crystals were observed in the sample of example 4 also at room temperature.

35

**Table 1** (comparison)

	Potassium sorbate % by weight	Room temperature	4 °C	-18 °C
Example 1	0.84	Clear solution	Clear solution	Crystals
Example 2	1.66	Clear solution	Crystals	Crystals
Example 3	3.27	Clear solution	Crystals	Crystals
Example 4	4.05	(Clear solution)	Crystals	Crystals

The result of example 1, which showed moderate preservability, 0.84 % by weight of potassium sorbate, equals about 0.6 % by weight of sorbic acid, this being too low a content for the intended purpose.

Examples 5 to 7.

50 g of potassium sorbate was dissolved into a 1 litre-volume of 99 % by weight formic acid, 99 % by weight propionic acid and 1:1 of both, 1 litre of each product containing sorbate in the same amount (examples 5 to 7).

The samples were treated as in the preceding example.

**Table 2**

	Acid	Room temperature	4 °C	-18 °C
Example 5 (comp.)	Propionic acid	Clear solution	Clear solution	Clear solution
Example 6 (comp.)	Formic acid	Clear solution	Crystals	Crystals
Example 7	Mixture of formic acid and propionic acid	Clear solution	Clear solution	Crystals

Examples 8 to 11.

Potassium sorbate was dissolved in mixtures of partly neutralised formic acid and propionic acid in the weight ratios shown in table 3 by heating the mixtures so as to yield clear solutions.

**Table 3.**

Compositions, % by weight	Example 8	Example 9	Example 10	Example 11
Formic acid	77.5	67.5	57.5	47.5
Propionic acid	20	30	40	50
Potassium sorbate	2.5	2.5	2.5	2.5
Neutralisation degree mol/100 mol	5.2	4.8	4.3	3.8
Water	14.3	12.5	10.6	8.8

All of the solutions remained clear at room temperature over the entire monitoring period, i.e. 3 weeks. The following table 4 shows the behaviour of three parallel samples in cooled solutions. The dash indicates a clear solution.

**Table 4**

Storage conditions	Example 8	Example 9	Example 10	Example 11
1 week at +1 °C	-	-	-	-
3 weeks at +1 °C	-	-	-	-

## Example 12.

10 g of potassium sorbate was dissolved in 100 g of mixtures of formic acid (99%) and propionic acid (99.5%) by heating so as to yield clear solutions. When cooled, the solutions developed deposits, which were removed by filtering at the end of a 1-week stabilisation period carried out at two different temperatures. The contents of sorbic acid, formic acid and propionic acid were analysed in the clear solutions. Table 5 indicates the sorbic acid result calculated as potassium sorbate and the percentage of propionic acid calculated on the sum of propionic and formic acid.

Table 5.

Sample		
Room temperature	Potassium sorbate % by weight	Amount % by weight of propionic acid of the sum of formic acid and propionic acid
1	2.3	0.0
2	3.4	9.9
3	4.8	20.2
4	6.4	30.2
5	7.7	40.1
6	9.7	50.1
Temperature 2 °C		
1	1.0	0.0
2	1.7	10.0
3	2.4	20.2
4	4.1	40.5
5	4.9	50.5
6	5.9	60.4
7	6.7	70.4
8	7.4	80.4

Figure 1 shows the solubility curves of potassium sorbate at room temperature 22 °C and at 2 °C. Compared to the solubility curve, the proportion of propionic acid



of the sum of formic acid and propionic acid should be about 5 to 10 percent units higher in practice.

Example 13.

5 The composition of the invention was needed in amounts of several tens of litres for swine tests. It was found that the dissolution of potassium sorbate as described above into a mixture of formic acid and propionic acid should be performed in a well equipped reactor, given highly evaporative acids that would require heating. As  
10 a result of the laboratory tests, a markedly easier manner of preparation was surprisingly found. In fact, it was found that preparation could be effected even at room temperature, with the potassium sorbate used in the composition first dissolved in propionic acid. The final composition was obtained by adding formic acid to a solution of potassium sorbate and propionic acid.

15 Example 14.

The composition of the invention was tested using weaned pigs and growing swine. The test comprised a total of 240 swine from 30 different litters. The composition of  
20 the invention or formic acid was added to the swine feed in a ratio of 8g/kg at the initial growth stage and in a ratio of 6g/kg at the growth and final growth stage. The composition of the invention was compared to a control without additives and to an antibiotic feed additive (Avilamycin) at the weaning and growth stage. Table 6 shows the test results with the administration of pig feed containing as additives  
25 avilamycin, formic acid and a mixture of potassium sorbate- formic acid-propionic acid. In the table A = avilamycin, F- formic acid and S = potassium sorbate and P = propionic acid.

**Table 6.**

Additive	Control	A	F	SFP
Weaning stage				
Initial weight/kg	7.5	7.5	7.6	7.8
Final weight/kg	17.2	19.3	18.5	19.0
Daily growth g/day	273	335	308	321
Growth and final growth stage				
Initial weight/kg	18.8	18.9	17.1	18.0
Final weight/kg	105.7	107.1	106.0	107.3
Daily growth g/day	834	845	869	880

- 5 The results indicate that swine grew more rapidly when the composition of the invention was added to the feed. The growth was also stronger than that of swine that received formic acid alone.

## Claims

1. An animal growth promoting composition, **characterised** in being a liquid solution comprising 1.1 to 5 % by weight of a sorbic acid or a sorbate calculated as sorbic acid, 45 to 93.9 % by weight of formic acid and 5 to 50 % by weight of propionic acid.
2. A composition as defined in claim 1, **characterised** in that the sorbate is potassium sorbate.
3. A composition as defined in claim 1 or 2, **characterised** in that the formic acid has a concentration of at least 70 % by weight.
4. A composition as defined in any of claims 1 to 3, **characterised** in that the formic acid is partly neutralised with ammonia.
5. A composition as defined in any of claims 1 to 4, **characterised** in that the composition additionally contains water.
6. A composition as defined in any of claims 1 to 5, **characterised** in that the composition comprises 2 to 4 % by weight of sorbic acid or a sorbate calculated as sorbic acid, 70 to 90 % by weight of partly neutralised formic acid and 8 to 20 % by weight of propionic acid and 0 to 20 % by weight of water.
7. A method for preparing an animal growth promoting composition, **characterised** in that sorbic acid or a sorbate is first dissolved in propionic acid, the solution being subsequently mixed into formic acid in order to form solution ready for use.
8. A method as defined in claim 7, **characterised** in that the sorbate is potassium sorbate.
9. A method as defined in claim 7, **characterised** in that the formic acid has a concentration of at least 70 % by weight.
10. A method as defined in claim 7, **characterised** in that the formic acid is partly neutralised with ammonia.

11. A method as defined in any of claims 7 to 10, **characterised** in that the solution ready for use comprises 2 to 5 % by weight of sorbic acid or a sorbate calculated as sorbic acid, 45 to 90 % by weight of formic acid and 8 to 50 % by weight of propionic acid.

5

12. A method as defined in claim 11, **characterised** in that the solution ready for use additionally contains water.

10

13. Use of a composition defined in claim 1 as a growth promoter for slaughter animals.

14. Use of a composition defined in claim 1 as a growth promoter, **characterised** in that the growth promoter is administered to animals in connection with feeding.


15

15. Use of a composition defined in claim 1 as a growth promoter, **characterised** in that the growth promoter is added to animal feed in liquid form or absorbed into a solid carrier.

20

16. Use of a composition defined in claim 15 as a growth promoter, **characterised** in that the animal feed is selected among organic products, such as feed raw materials, feed preparations, corn, vegetable and animal protein feed.

FIGURE 1. Solubility curves of potassium sorbate at different temperatures



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/IT 03/00504

## A. CLASSIFICATION OF SUBJECT MATTER

IPC7: A23K 1/16

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: A23K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-INTERNAL WPI DATA, CAPLUS, FSTA, CABA, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Livestock Production Science, Volume 73, 2002, Kirsi Partanen et al, "Performance of growing-finishing pigs fed medium- or high-fibre diets supplement with avilamycin, formic acid or formic acid - sorbate blend", pages 139-152, --	1-16
A	EP 1205115 A2 (NUTRINOVA NUTRITION SPECIALITES & FOOD INGREDIENTS GMBH), 15 May 2002 (15.05.02) --	1-16
A	US 4083999 A (DRURY ET AL), 11 April 1978 (11.04.78) --	1-16

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"B" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

30 Sept 2003

Date of mailing of the international search report

08-10-2003

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

DAGMAR JÄRVMAN/BS

Telephone No. +46 8 782 25 00

## INTERNATIONAL SEARCH REPORT

Internal application No.

PCT/PI 03/00504

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0701778 A2 (HOECHST AKTIENGESELLSCHAFT), 20 March 1996 (20.03.96)  --	1-16
A	GB 1452169 A (FEED SERVICE (LIVESTOCK) LIMITED), 13 October 1976 (13.10.76)  -- -----	1-16

## INTERNATIONAL SEARCH REPORT

Information on patent family members

06/09/03

International application No.

PCT/PT 03/00504

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
EP	1205115	A2	15/05/02	AU	8933901 A	16/05/02
				DE	10055390 A	29/05/02
				JP	2002191293 A	09/07/02
				NZ	514923 A	25/07/03
				US	2002086090 A	04/07/02
-----						
US	4083999	A	11/04/78	NONE		
-----						
EP	0701778	A2	20/03/96	BR	9504056 A	24/09/96
				CN	1131514 A	25/09/96
				CZ	9502424 A	17/04/96
				DE	19510158 A	28/03/96
				DE	29521537 U	30/04/98
				FI	954362 A	20/03/96
				JP	8173058 A	09/07/96
				US	5888569 A	30/03/99
-----						
GB	1452169	A	13/10/76	FR	2267710 A,B	14/11/75
-----						



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**